

Lewenstein-Sanpera Decomposition for Iso-concurrence Decomposable States

S. J. Akhtarshenas^{a,b,c} *, M. A. Jafarizadeh^{a,b,c} †

^aDepartment of Theoretical Physics and Astrophysics, Tabriz University, Tabriz 51664, Iran.

^bInstitute for Studies in Theoretical Physics and Mathematics, Tehran 19395-1795, Iran.

^cResearch Institute for Fundamental Sciences, Tabriz 51664, Iran.

February 9, 2008

*E-mail:akhtarshenas@tabrizu.ac.ir

†E-mail:jafarizadeh@tabrizu.ac.ir

Abstract

We obtain Lewenstein-Sanpera decomposition of iso-concurrence decomposable states of $2 \otimes 2$ quantum systems. It is shown that in these systems average concurrence of the decomposition is equal to the concurrence of the state and also it is equal to the amount of violation of positive partial transpose criterion. It is also shown that the product states introduced by Wootters in [W. K. Wootters, Phys. Rev. Lett. **80** 2245 (1998)] form the best separable approximation ensemble for these states.

Keywords: Quantum entanglement, Iso-concurrence decomposable states, Lewenstein-Sanpera decomposition, Concurrence

PACs Index: 03.65.Ud

1 Introduction

During the past decade an increasing study has been made on entanglement, although it was discovered many decades ago by Einstein and Schrödinger [1, 2]. This is because of potential resource that entanglement provides for quantum communication and information processing [3, 4, 5]. Entanglement usually arises from quantum correlations between separated subsystems which can not be created by local actions on each subsystems. By definition, a bipartite mixed state ρ is said to be separable if it can be expressed as

$$\rho = \sum_i w_i \rho_i^{(1)} \otimes \rho_i^{(2)}, \quad w_i \geq 0, \quad \sum_i w_i = 1,$$

where $\rho_i^{(1)}$ and $\rho_i^{(2)}$ denote density matrices of subsystems 1 and 2 respectively. Otherwise the state is entangled.

The central tasks of quantum information theory is to characterize and quantify entangled states. A first attempt in characterization of entangled states has been made by Peres and Horodecki family [6, 7]. It was shown that a necessary condition for separability of a two partite system is that its partial transposition be positive. Horodeckis showed that this condition is sufficient for separability of composite systems only for dimensions $2 \otimes 2$ and $2 \otimes 3$.

There is also an increasing attention in quantifying entanglement, particularly for mixed states of a bipartite system, and a number of measures have been proposed [5, 8, 9, 10]. Among them the entanglement of formation has more importance, since it intends to quantify the resources needed to create a given entangled state.

An interesting description of entanglement is Lewenstein-Sanpera decomposition [11]. Lewenstein and Sanpera in [11] have shown that any two partite density matrix can be represented optimally as a sum of a separable state (called best separable approximation (BSA)) and an entan-

gled state. They have also shown that for 2-qubit systems the decomposition reduces to a mixture of a mixed separable state and an entangled pure state, thus all non-separability content of the state is concentrated in the pure entangled state. This leads to an unambiguous measure of entanglement for any 2-qubit state as entanglement of pure state multiplied by the weight of pure part in the decomposition.

In the Ref. [11], the numerical method for finding the BSA has been reported. Also in $2 \otimes 2$ systems some analytical results for special states were found in [12]. An analytical expression for L-S decomposition of Bell decomposable states and a class of states obtained from them via some LQCC action is also obtained in [13].

In this paper we introduce iso-concurrence decomposable (ICD) states. We give an analytical expression for L-S decomposition of these states to show that the average concurrence of the decomposition is equal to their concurrence in these states. As a byproduct we also show that the concurrence of the considered states is equal to amount of violation of positive partial transpose criterion. It is also shown that the product states introduced by Wootters in [10] form BSA ensemble for these states.

Considering a real matrix parameterization of density matrices, Verstraete et al. in [14, 15] showed that an local quantum operations and classical communications (LQCC) corresponds to left and right multiplication by a Lorentz matrix. They also have shown that the real parameterization matrices $R_{ij} = Tr(\rho \sigma_i \sigma_j)$ can be decomposed as $R = L_1 \Sigma L_2^T$, where L_1 and L_2^T are proper Lorentz transformations and Σ has either a diagonal form correspond to Bell decomposable states or it has a non-diagonal form. For the case that the matrix Σ becomes Bell decomposable state we presented optimal decomposition for states obtained from them via some LQCC actions [13]. In other cases that Σ has non-diagonal form, corresponding matrix is a special cases of ICD states,

where we consider it in this paper, and investigation of action of LQCC on these states will appear in another paper.

The paper is organized as follows. In section 2 we introduce ICD states, then we study their separability properties. The concurrence of these states is evaluated in section 3, via the method presented by Wootters in [10]. In section 4 we obtain L-S decomposition of these states and we prove that thus obtained decomposition is optimal. The paper is ended with a brief conclusion in section 5.

2 Iso-concurrence decomposable states

In this section we define iso-concurrence decomposable (ICD) states, then we give their separability condition. The iso-concurrence states are defined by

$$|\psi_1\rangle = \cos \theta |\uparrow\uparrow\rangle + \sin \theta |\downarrow\downarrow\rangle, \quad (2-1)$$

$$|\psi_2\rangle = \sin \theta |\uparrow\uparrow\rangle - \cos \theta |\downarrow\downarrow\rangle, \quad (2-2)$$

$$|\psi_3\rangle = \cos \theta |\uparrow\downarrow\rangle + \sin \theta |\downarrow\uparrow\rangle, \quad (2-3)$$

$$|\psi_4\rangle = \sin \theta |\uparrow\downarrow\rangle - \cos \theta |\downarrow\uparrow\rangle. \quad (2-4)$$

It is quite easy to see that the above states are orthogonal and thus they span the Hilbert space of $2 \otimes 2$ systems. Also by choosing $\theta = \frac{\pi}{4}$ the above states reduce to Bell states. Now we can define ICD states as

$$\rho = \sum_{i=1}^4 p_i |\psi_i\rangle \langle \psi_i|, \quad 0 \leq p_i \leq 1, \quad \sum_{i=1}^4 p_i = 1. \quad (2-5)$$

These states form a four simplex (tetrahedral) with its vertices defined by $p_1 = 1, p_2 = 1, p_3 = 1$ and $p_4 = 1$, respectively.

A necessary condition for separability of composite quantum systems is presented by Peres [6]. He showed that if a state is separable then the matrix obtained from partial transposition must be positive. Horodecki family [7] have shown that Peres criterion provides sufficient condition only for separability of mixed quantum states of dimensions $2 \otimes 2$ and $2 \otimes 3$. This implies that the state given in Eq. (2-5) is separable if and only if the following inequalities are satisfied

$$(p_1 - p_2) \leq \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta}, \quad (2-6)$$

$$(p_2 - p_1) \leq \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta}, \quad (2-7)$$

$$(p_3 - p_4) \leq \sqrt{4p_1p_2 + (p_1 - p_2)^2 \sin^2 2\theta}, \quad (2-8)$$

$$(p_4 - p_3) \leq \sqrt{4p_1p_2 + (p_1 - p_2)^2 \sin^2 2\theta}. \quad (2-9)$$

Inequalities (2-6) to (2-9) divide tetrahedral of density matrices to five regions. Central regions, defined by the above inequalities, form a deformed octahedral and are separable states. In four other regions one of the above inequality will not hold, therefore they represent entangled states. In the next sections we consider entangled states corresponding to violation of inequality (2-6) i.e. the states which satisfy the following inequality

$$(p_1 - p_2) > \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta}. \quad (2-10)$$

All other ICD states can be obtain via local unitary transformations.

3 Concurrence

In this section we first review concurrence of mixed states. From the various measures proposed to quantify entanglement, the entanglement of formation has a special position which in fact intends to quantify the resources needed to create a given entangled state [5]. Wootters in [10] has shown

that for a 2-qubit system entanglement of formation of a mixed state ρ can be defined as

$$E(\rho) = H\left(\frac{1}{2} + \frac{1}{2}\sqrt{1 - C^2}\right), \quad (3-11)$$

where $H(x) = -x \ln x - (1 - x) \ln (1 - x)$ is the binary entropy and the concurrence $C(\rho)$ is defined by

$$C(\rho) = \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\}, \quad (3-12)$$

where the λ_i are the non-negative eigenvalues, with λ_1 being the largest one, of the Hermitian matrix $R \equiv \sqrt{\sqrt{\rho}\tilde{\rho}\sqrt{\rho}}$ and

$$\tilde{\rho} = (\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y), \quad (3-13)$$

where ρ^* is the complex conjugate of ρ when it is expressed in a fixed basis such as $\{|\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle\}, \{|\downarrow\uparrow\rangle, |\downarrow\downarrow\rangle\}$ and σ_y represent Pauli matrix in local basis $\{|\uparrow\rangle, |\downarrow\rangle\}$.

In order to obtain the concurrence of ICD states we the follow method presented by Wootters in [10]. Starting from spectral decomposition for ICD states given in Eq. (2-5), we define subnormalized orthogonal eigenvectors $|v_i\rangle$ as

$$|v_i\rangle = \sqrt{p_i} |\psi_i\rangle, \quad \langle v_i | v_j \rangle = p_i \delta_{ij}. \quad (3-14)$$

Now we can define states $|x_i\rangle$ by

$$|x_i\rangle = \sum_j^4 U_{ij}^* |v_i\rangle, \quad \text{for } i = 1, 2, 3, 4, \quad (3-15)$$

such that

$$\langle x_i | \tilde{x}_j \rangle = (U\tau U^T)_{ij} = \lambda_i \delta_{ij}, \quad (3-16)$$

where $\tau_{ij} = \langle v_i | v_j \rangle$ is a symmetric but not necessarily Hermitian matrix. To construct $|x_i\rangle$ we consider the fact that for any symmetric matrix τ one can always find a unitary matrix U in such

a way that λ_i are real and non-negative, that is, they are the square roots of eigenvalues of $\tau\tau^*$ which are same as eigenvalues of R . Moreover one can always find U such that λ_1 being the largest one.

By using the above protocol we get the following expression for state of ρ given in Eq. (2-5)

$$\tau = \begin{pmatrix} -p_1 \sin 2\theta & \sqrt{p_1 p_2} \cos 2\theta & 0 & 0 \\ \sqrt{p_1 p_2} \cos 2\theta & p_2 \sin 2\theta & 0 & 0 \\ 0 & 0 & p_3 \sin 2\theta & -\sqrt{p_3 p_4} \cos 2\theta \\ 0 & 0 & -\sqrt{p_3 p_4} \cos 2\theta & -p_4 \sin 2\theta \end{pmatrix}. \quad (3-17)$$

Now it is easy to evaluate λ_i which yields

$$\begin{aligned} \lambda_1 &= \frac{1}{2} \left((p_1 - p_2) \sin 2\theta + \sqrt{4p_1 p_2 + (p_1 - p_2)^2 \sin^2 2\theta} \right), \\ \lambda_2 &= \frac{1}{2} \left((p_2 - p_1) \sin 2\theta + \sqrt{4p_1 p_2 + (p_1 - p_2)^2 \sin^2 2\theta} \right), \\ \lambda_3 &= \frac{1}{2} \left((p_3 - p_4) \sin 2\theta + \sqrt{4p_3 p_4 + (p_3 - p_4)^2 \sin^2 2\theta} \right), \\ \lambda_4 &= \frac{1}{2} \left((p_4 - p_3) \sin 2\theta + \sqrt{4p_3 p_4 + (p_3 - p_4)^2 \sin^2 2\theta} \right). \end{aligned} \quad (3-18)$$

Thus one can evaluate the concurrence of ICD states as

$$C = (p_1 - p_2) \sin 2\theta - \sqrt{4p_3 p_4 + (p_3 - p_4)^2 \sin^2 2\theta}. \quad (3-19)$$

It is worth to note that thus obtained concurrence is equal to the amount of violation of inequality (2-10).

Finally we introduce the unitary matrix U which is going to be used later

$$U = \begin{pmatrix} i\alpha_1 & -i\alpha_2 & 0 & 0 \\ \alpha_2 & \alpha_1 & 0 & 0 \\ 0 & 0 & \alpha_3 & -\alpha_4 \\ 0 & 0 & i\alpha_4 & i\alpha_3 \end{pmatrix}, \quad (3-20)$$

where

$$\begin{aligned}
\alpha_1 &= \frac{\left((p_1 + p_2) \sin 2\theta + \sqrt{4p_1p_2 + (p_1 - p_2)^2 \sin^2 2\theta} \right)}{\sqrt{2} \left(4p_1p_2 \cos^2 2\theta + (p_1 + p_2)^2 \sin^2 2\theta + (p_1 + p_2) \sin 2\theta \sqrt{4p_1p_2 + (p_1 - p_2)^2 \sin^2 2\theta} \right)}, \\
\alpha_2 &= \frac{\sqrt{2p_1p_2} \cos 2\theta}{\left(4p_1p_2 \cos^2 2\theta + (p_1 + p_2)^2 \sin^2 2\theta + (p_1 + p_2) \sin 2\theta \sqrt{4p_1p_2 + (p_1 - p_2)^2 \sin^2 2\theta} \right)}, \\
\alpha_3 &= \frac{\left((p_3 + p_4) \sin 2\theta + \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta} \right)}{\sqrt{2} \left(4p_3p_4 \cos^2 2\theta + (p_3 + p_4)^2 \sin^2 2\theta + (p_3 + p_4) \sin 2\theta \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta} \right)}, \\
\alpha_4 &= \frac{\sqrt{2p_3p_4} \cos 2\theta}{\left(4p_3p_4 \cos^2 2\theta + (p_3 + p_4)^2 \sin^2 2\theta + (p_3 + p_4) \sin 2\theta \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta} \right)}.
\end{aligned} \tag{3-21}$$

4 Lewenstein-Sanpera decomposition

According to Lewenstein-Sanpera decomposition [11], any 2-qubit density matrix ρ can be written as

$$\rho = \lambda \rho_{sep} + (1 - \lambda) |\psi\rangle \langle \psi|, \quad \lambda \in [0, 1], \tag{4-22}$$

where ρ_{sep} is a separable density matrix and $|\psi\rangle$ is a pure entangled state. The Lewenstein-Sanpera decomposition of a given density matrix ρ is not unique and, in general, there is a continuum set of L-S decomposition to choose from. The optimal decomposition is, however, unique for which λ is maximal and

$$\rho = \lambda^{(opt)} \rho_{sep}^{(opt)} + (1 - \lambda^{(opt)}) |\psi^{(opt)}\rangle \langle \psi^{(opt)}|, \quad \lambda^{(opt)} \in [0, 1]. \tag{4-23}$$

Lewenstein and Sanpera in [11] have shown that any other decomposition of the form $\rho = \tilde{\lambda} \tilde{\rho}_{sep} + (1 - \tilde{\lambda}) |\tilde{\psi}\rangle \langle \tilde{\psi}|$ with $\tilde{\rho} \neq \rho^{(opt)}$ necessarily implies that $\tilde{\lambda} < \lambda^{(opt)}$ [11]. We refer to Eq. (4-23) as

optimal decomposition in the sense that λ is maximal and ρ_s is the best separable approximation (BSA).

Here in this section we would like to obtain L-S decomposition for ICD states. Let us consider entangled state ρ belonging to entangled region defined by Eq. (2-10). We start by writing density matrix ρ as a convex sum of pure state $|\psi_1\rangle$ and separable state ρ_s as

$$\rho = \lambda \rho_s + (1 - \lambda) |\psi_1\rangle \langle \psi_1|. \quad (4-24)$$

where, ρ_s corresponds to intersection of the line passing through the points corresponding to $|\psi_1\rangle \langle \psi_1|$ and ρ with 2-dimensional manifold defined by the equality in relation (2-6). Expanding separable state ρ_s as $\rho_s = \sum_{i=1}^4 p'_i |\psi_i\rangle \langle \psi_i|$ and using Eq. (2-5) for ρ we arrive at the following results

$$p'_1 = \frac{p_1 + \lambda - 1}{\lambda}, \quad p'_i = \frac{p_i}{\lambda} \quad \text{for } i = 2, 3, 4, \quad (4-25)$$

where

$$\lambda = 1 - p_1 + p_2 + \sqrt{\frac{4p_3p_4}{\sin^2 2\theta} + (p_3 - p_4)^2}. \quad (4-26)$$

Now using the fact that for pure states defined in Eqs. (2-1) to (2-4) we have $C(|\psi_i\rangle) = \sin 2\theta$, one can easily evaluate average concurrence of the decomposition given in Eq. (4-24) as

$$(1 - \lambda)C(|\psi_1\rangle) = (p_1 - p_2) \sin 2\theta - \sqrt{4p_3p_4 + (p_3 - p_4)^2 \sin^2 2\theta}. \quad (4-27)$$

Equation (4-27) shows that average concurrence of ρ in the L-S decomposition given in (4-24) is equal to its concurrence. As the concurrence of a mixed state is defined as the minimum of the average concurrence over all decompositions of the state in terms of pure states [10], so according to Eq. (4-24) we should have $C(\rho) \leq (1 - \lambda)C(\psi)$. But Eq. (4-27) shows that the optimal decomposition of ICD states saturate this inequality.

In the rest of this section we will prove that the decomposition (4-24) is the optimal one. To do so we have to find a decomposition for ρ_s in terms of product states $|e_\alpha, f_\alpha\rangle$, i.e.

$$\rho_s = \sum_{\alpha} \Lambda_{\alpha} |e_{\alpha}, f_{\alpha}\rangle \langle e_{\alpha}, f_{\alpha}| \quad (4-28)$$

such that the following conditions are satisfied [11]

i) All Λ_{α} are maximal with respect to $\rho_{\alpha} = \Lambda_{\alpha} |e_{\alpha}, f_{\alpha}\rangle \langle e_{\alpha}, f_{\alpha}| + (1 - \lambda) |\psi_1\rangle \langle \psi_1|$ and projector $P_{\alpha} = |e_{\alpha}, f_{\alpha}\rangle \langle e_{\alpha}, f_{\alpha}|$.

ii) All pairs $(\Lambda_{\alpha}, \Lambda_{\beta})$ are maximal with respect to $\rho_{\alpha\beta} = \Lambda_{\alpha} |e_{\alpha}, f_{\alpha}\rangle \langle e_{\alpha}, f_{\alpha}| + \Lambda_{\beta} |e_{\beta}, f_{\beta}\rangle \langle e_{\beta}, f_{\beta}| + (1 - \lambda) |\psi_1\rangle \langle \psi_1|$ and the pairs of projector (P_{α}, P_{β}) .

Then according to [11] ρ_s is BSA and the decomposition given in Eq. (4-24) is optimal.

Lewenstein and Sanpera in [11] have shown that Λ is maximal with respect to ρ and $P = |\psi\rangle \langle \psi|$ iff a) if $|\psi\rangle \notin \mathcal{R}(\rho)$ then $\Lambda = 0$, and b) if $|\psi\rangle \in \mathcal{R}(\rho)$ then $\Lambda = \langle \psi | \rho^{-1} | \psi \rangle^{-1} > 0$. They have also shown that a pair (Λ_1, Λ_2) is maximal with respect to ρ and a pair of projectors (P_1, P_2) iff: a) if $|\psi_1\rangle, |\psi_2\rangle$ do not belong to $\mathcal{R}(\rho)$ then $\Lambda_1 = \Lambda_2 = 0$; b) if $|\psi_1\rangle$ does not belong, while $|\psi_2\rangle \in \mathcal{R}(\rho)$ then $\Lambda_1 = 0, \Lambda_2 = \langle \psi_2 | \rho^{-1} | \psi_2 \rangle^{-1}$; c) if $|\psi_1\rangle, |\psi_2\rangle \in \mathcal{R}(\rho)$ and $\langle \psi_1 | \rho^{-1} | \psi_2 \rangle = 0$ then $\Lambda_i = \langle \psi_i | \rho^{-1} | \psi_i \rangle^{-1}$, $i = 1, 2$; d) finally, if $|\psi_1\rangle, |\psi_2\rangle \in \mathcal{R}(\rho)$ and $\langle \psi_1 | \rho^{-1} | \psi_2 \rangle \neq 0$ then

$$\begin{aligned} \Lambda_1 &= (\langle \psi_2 | \rho^{-1} | \psi_2 \rangle - |\langle \psi_1 | \rho^{-1} | \psi_2 \rangle|) / D, \\ \Lambda_2 &= (\langle \psi_1 | \rho^{-1} | \psi_1 \rangle - |\langle \psi_1 | \rho^{-1} | \psi_2 \rangle|) / D, \end{aligned} \quad (4-29)$$

where $D = \langle \psi_1 | \rho^{-1} | \psi_1 \rangle \langle \psi_2 | \rho^{-1} | \psi_2 \rangle - |\langle \psi_1 | \rho^{-1} | \psi_2 \rangle|^2$.

Now we return to show that the decomposition given in (4-24) is optimal. Wootters in [10] has shown that any $2 \otimes 2$ separable density matrix can be expanded in terms of following product states

$$|z_1\rangle = \frac{1}{2} \left(e^{i\theta_1} |x_1\rangle + e^{i\theta_2} |x_2\rangle + e^{i\theta_3} |x_3\rangle + e^{i\theta_4} |x_4\rangle \right), \quad (4-30)$$

$$|z_2\rangle = \frac{1}{2} \left(e^{i\theta_1} |x_1\rangle + e^{i\theta_2} |x_2\rangle - e^{i\theta_3} |x_3\rangle - e^{i\theta_4} |x_4\rangle \right), \quad (4-31)$$

$$|z_3\rangle = \frac{1}{2} \left(e^{i\theta_1} |x_1\rangle - e^{i\theta_2} |x_2\rangle + e^{i\theta_3} |x_3\rangle - e^{i\theta_4} |x_4\rangle \right), \quad (4-32)$$

$$|z_4\rangle = \frac{1}{2} \left(e^{i\theta_1} |x_1\rangle - e^{i\theta_2} |x_2\rangle - e^{i\theta_3} |x_3\rangle + e^{i\theta_4} |x_4\rangle \right), \quad (4-33)$$

provided that $\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4 \leq 0$. Now, the zero concurrence is guaranteed by choosing phases θ_i , $i = 1, 2, 3, 4$ satisfying the relation $\sum_{j=1} e^{2i\theta_j} \lambda_j = 0$.

Now using the fact that for marginal separable states ρ_s (located at the boundary of separable region) the eigenvalues λ_i satisfy the constraint $\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4 = 0$, we can choose the phase factors θ_i as $\theta_2 = \theta_3 = \theta_4 = \theta_1 + \frac{\pi}{2}$. Choosing $\theta_1 = 0$ we arrive at the following product ensemble for ρ_s

$$\begin{aligned} |z_1\rangle &= \frac{1}{2} (-i(\alpha'_1 + \alpha'_2) \sqrt{p'_1} |\psi_1\rangle - i(\alpha'_1 - \alpha'_2) \sqrt{p'_2} |\psi_2\rangle \\ &\quad - (\alpha'_4 + i\alpha'_3) \sqrt{p'_3} |\psi_3\rangle - (\alpha'_3 - i\alpha'_4) \sqrt{p'_4} |\psi_4\rangle), \\ |z_2\rangle &= \frac{1}{2} (-i(\alpha'_1 + \alpha'_2) \sqrt{p'_1} |\psi_1\rangle - i(\alpha'_1 - \alpha'_2) \sqrt{p'_2} |\psi_2\rangle \\ &\quad + (\alpha'_4 + i\alpha'_3) \sqrt{p'_3} |\psi_3\rangle + (\alpha'_3 - i\alpha'_4) \sqrt{p'_4} |\psi_4\rangle), \\ |z_3\rangle &= \frac{1}{2} (-i(\alpha'_1 - \alpha'_2) \sqrt{p'_1} |\psi_1\rangle + i(\alpha'_1 + \alpha'_2) \sqrt{p'_2} |\psi_2\rangle \\ &\quad + (\alpha'_4 - i\alpha'_3) \sqrt{p'_3} |\psi_3\rangle + (\alpha'_3 + i\alpha'_4) \sqrt{p'_4} |\psi_4\rangle), \\ |z_4\rangle &= \frac{1}{2} (-i(\alpha'_1 - \alpha'_2) \sqrt{p'_1} |\psi_1\rangle + i(\alpha'_1 + \alpha'_2) \sqrt{p'_2} |\psi_2\rangle \\ &\quad - (\alpha'_4 - i\alpha'_3) \sqrt{p'_3} |\psi_3\rangle - (\alpha'_3 + i\alpha'_4) \sqrt{p'_4} |\psi_4\rangle), \end{aligned} \quad (4-34)$$

where α'_i are defined in the same form as given in Eq. (3-21) provided that they are expressed in terms of p'_i (coordinates of ρ_s).

Now in order to prove that the decomposition given in Eq. (4-22) is optimal we first consider the cases that ρ has full rank. Let us consider the set of four product vectors $\{|z_i\rangle\}$ and one entangled state $|\psi_1\rangle$. In Ref. [10] it is shown that the ensemble $\{|z_i\rangle\}$ are linearly independent, also it is

straightforward to see that three vectors $|z_\alpha\rangle$, $|z_\beta\rangle$ and $|\psi_1\rangle$ are linearly independent. Now let us consider matrices $\rho_\alpha = \Lambda_\alpha |z_\alpha\rangle \langle z_\alpha| + (1-\lambda) |\psi_1\rangle \langle \psi_1|$. Due to independence of $|z_\alpha\rangle$ and $|\psi_1\rangle$ we can deduce that the range of ρ_α is two dimensional, thus after restriction to its range and defining dual basis $|\hat{z}_\alpha\rangle$ and $|\hat{\psi}_1\rangle$ we can expand restricted inverse ρ_α^{-1} as $\rho_\alpha^{-1} = \Lambda_\alpha^{-1} |\hat{z}_\alpha\rangle \langle \hat{z}_\alpha| + (1-\lambda)^{-1} |\hat{\psi}_1\rangle \langle \hat{\psi}_1|$ (see appendix). Using Eq. (5-35) it is easy to see that $\langle z_\alpha | \rho_\alpha^{-1} | z_\alpha \rangle = \Lambda_\alpha^{-1}$. This shows that Λ_α are maximal with respect to ρ_α and the projector $P_\alpha = |z_\alpha\rangle \langle z_\alpha|$.

Similarly considering matrices $\rho_{\alpha\beta} = \Lambda_\alpha |z_\alpha\rangle \langle z_\alpha| + \Lambda_\beta |z_\beta\rangle \langle z_\beta| + (1-\lambda) |\psi_1\rangle \langle \psi_1|$ and considering the independence of vectors $|z_\alpha\rangle$, $|z_\beta\rangle$ and $|\psi_1\rangle$ we see that rang of $\rho_{\alpha\beta}$ is three dimensional where after restriction to its range and defining their dual basis $|\hat{z}_\alpha\rangle$, $|\hat{z}_\beta\rangle$ and $|\hat{\psi}_1\rangle$ we can write restricted inverse $\rho_{\alpha\beta}^{-1}$ as $\rho_{\alpha\beta}^{-1} = \Lambda_\alpha^{-1} |\hat{z}_\alpha\rangle \langle \hat{z}_\alpha| + \Lambda_\beta^{-1} |\hat{z}_\beta\rangle \langle \hat{z}_\beta| + (1-\lambda)^{-1} |\hat{\psi}_1\rangle \langle \hat{\psi}_1|$. Then it is straightforward to get $\langle \hat{z}_\alpha | \rho_{\alpha\beta}^{-1} | \hat{z}_\alpha \rangle = \Lambda_\alpha^{-1}$, $\langle \hat{z}_\beta | \rho_{\alpha\beta}^{-1} | \hat{z}_\beta \rangle = \Lambda_\beta^{-1}$ and $\langle \hat{z}_\alpha | \rho_{\alpha\beta}^{-1} | \hat{z}_\beta \rangle = 0$. This implies that the pair $(\Lambda_\alpha, \Lambda_\beta)$ are maximal with respect to $\rho_{\alpha\beta}$ and the pair of projectors (P_α, P_β) , thus complete the proof that the decomposition given in Eq. (4-22) is optimal.

We now consider the cases that ρ has not full rank. First let us consider $p'_2 = 0$ case. In this case the pairs $\{|z_1\rangle, |z_2\rangle\}$ and also $\{|z_3\rangle, |z_4\rangle\}$ are no longer independent with respect to $|\psi_1\rangle$. In this cases we can easily evaluate $|\psi_1\rangle$ in terms of $|z_1\rangle$ and $|z_2\rangle$ then matrix ρ_{12} can be written in terms of two basis $|z_1\rangle$ and $|z_2\rangle$ which yields after some calculations, $\langle z_1 | \rho_{12}^{-1} | z_1 \rangle = \frac{\Lambda_2(\alpha'_1 + \alpha'_2)^2 p'_1 + (1-\lambda)}{\Gamma_{12}}$, $\langle z_2 | \rho_{12}^{-1} | z_2 \rangle = \frac{\Lambda_1(\alpha'_1 + \alpha'_2)^2 p'_1 + (1-\lambda)}{\Gamma_{12}}$ and $\langle z_1 | \rho_{12}^{-1} | z_2 \rangle = \frac{-(1-\lambda)}{\Gamma_{12}}$ where $\Gamma_{12} = \Lambda_1 \Lambda_2 (\alpha'_1 + \alpha'_2)^2 p'_1 + (1-\lambda)(\Lambda_1 + \Lambda_2)$. Using the above results together with Eqs. (4-29) we obtain the maximality of pair (Λ_1, Λ_2) with respect to ρ_{12} and the pair of projectors (P_1, P_2) . Similarly we can express $|\psi_1\rangle$ in terms of $|z_3\rangle$ and $|z_4\rangle$ and evaluate ρ_{34}^{-1} , which get $\langle z_3 | \rho_{34}^{-1} | z_3 \rangle = \frac{\Lambda_4(\alpha'_1 - \alpha'_2)^2 p'_1 + (1-\lambda)}{\Gamma_{34}}$, $\langle z_4 | \rho_{34}^{-1} | z_4 \rangle = \frac{\Lambda_3(\alpha'_1 - \alpha'_2)^2 p'_1 + (1-\lambda)}{\Gamma_{34}}$ and $\langle z_3 | \rho_{34}^{-1} | z_4 \rangle = \frac{-(1-\lambda)}{\Gamma_{34}}$ with $\Gamma_{34} = \Lambda_3 \Lambda_4 (\alpha'_1 - \alpha'_2)^2 p'_1 + (1-\lambda)(\Lambda_3 + \Lambda_4)$, together with Eqs. (4-29) we obtain the maximality of pair (Λ_3, Λ_4) with respect to ρ_{34} and the

pair of projectors (P_3, P_4) . For other choices of α and β three vectors $|z_\alpha\rangle, |z_\beta\rangle$ and $|\psi_1\rangle$ remain linearly independent, thus with the above mentioned method we can prove the maximality of pairs $(\Lambda_\alpha, \Lambda_\beta)$.

Also in the cases of $p'_3 = 0$ or $p'_4 = 0$, three vectors $|z_\alpha\rangle, |z_\beta\rangle$ and $|\psi_1\rangle$ remain linearly independent thus maximality of pairs $(\Lambda_\alpha, \Lambda_\beta)$ can be proven in the same method of full rank one.

Now let us consider the cases that rank ρ is 2. In this case we have to write the separable part in terms of rank one projection operators which are not product any more but as far as the maximality of pairs (the requirement for the maximality of separable part of density matrix in this case) are concerned the separability of the projectors does not play any role.

For $p_2 = p_3 = 0$ we have $p'_1 = p'_4 = \frac{1}{2}$, therefore ρ_s can be written as

$$\rho_s = \frac{1}{2} (|\psi_1\rangle \langle\psi_1| + |\psi_4\rangle \langle\psi_4|) = \frac{1}{2} (|e_1, f_1\rangle \langle e_1, f_1| + |e_2, f_2\rangle \langle e_2, f_2|),$$

where $|e_{1,2}, f_{1,2}\rangle = (\cos \theta |\uparrow\rangle \mp i \sin \theta |\downarrow\rangle) \otimes \frac{1}{\sqrt{2}} (|\uparrow\rangle \pm i |\downarrow\rangle)$. Now by writing $|\psi_1\rangle$ in terms of product states $|e_1, f_1\rangle$ and $|e_2, f_2\rangle$ the matrix ρ_{12} can be expressed in terms of two basis $|e_1, f_1\rangle, |e_2, f_2\rangle$ and we get $\langle e_1, f_1 | \rho_{12}^{-1} | e_1, f_1 \rangle = \frac{2\Lambda_2 + (1-\lambda)}{\Gamma_{12}}$, $\langle e_2, f_2 | \rho_{12}^{-1} | e_2, f_2 \rangle = \frac{2\Lambda_1 + (1-\lambda)}{\Gamma_{12}}$ and $\langle e_1, f_1 | \rho_{12}^{-1} | e_2, f_2 \rangle = \frac{-(1-\lambda)}{\Gamma_{12}}$ where $\Gamma_{12} = 2\Lambda_1\Lambda_2 + (1-\lambda)(\Lambda_1 + \Lambda_2)$. Using the above results together with Eqs. (4-29) we obtain the maximality of pair (Λ_1, Λ_2) with respect to ρ_{12} and the pair of projectors (P_1, P_2) . The other rank-2 cases can be proved via similar method.

5 Conclusion

We have derived Lewenstein-Sanpera decomposition for iso-concurrence decomposable states. It is shown that for these states the average concurrence of L-S decomposition is equal to the concurrence of the states. It is also shown that Wootters product states defined in [10] form the best separable

approximation ensemble for these states.

Appendix

Let us consider the set of linearly independent vectors $\{|\phi_i\rangle\}$, then one can define their dual vectors $\{|\hat{\phi}_i\rangle\}$ such that the following relation

$$\langle \hat{\phi}_i | \phi_j \rangle = \delta_{ij} \quad (5-35)$$

hold. It is straightforward to show that the $\{|\phi_i\rangle\}$ and their dual $\{|\hat{\phi}_i\rangle\}$ posses the following completeness relation

$$\sum_i |\hat{\phi}_i\rangle \langle \phi_i| = I, \quad \sum_i |\phi_i\rangle \langle \hat{\phi}_i| = I. \quad (5-36)$$

Consider an invertible operator M which is expanded in terms of states $|\phi_i\rangle$ as

$$M = \sum_i a_{ij} |\phi_i\rangle \langle \phi_j| \quad (5-37)$$

Then the inverse of M denoted by M^{-1} can be expanded in terms of dual bases as

$$M^{-1} = \sum_i b_{ij} |\hat{\phi}_i\rangle \langle \hat{\phi}_j| \quad (5-38)$$

where $b_{ij} = (A^{-1})_{ij}$ and $A_{ij} = a_{ij}$.

References

- [1] A. EINSTEIN, B. PODOLSKY AND ROSEN, *Phys. Rev.* **47**, 777 (1935).
- [2] E. SCHRÖDINGER, *Naturwissenschaften.* **23** 807 (1935).
- [3] C. H. BENNETT, AND S. J. WIESNER, *Phys. Rev. Lett.* **69**, 2881 (1992).
- [4] C. H. BENNETT, G. BRASSARD, C. CRÉPEAU, R. JOZSA, A PERES AND W. K. WOOTTERS, *Phys. Rev. Lett.* **70**, 1895 (1993).

- [5] C. H. BENNETT, D. P. DiVINCENZO, J. A. SMOLIN AND W.K. WOOTTERS, *Phys. Rev. A* **54**, 3824 (1996).
- [6] A. PERES, *Phys. Rev. Lett.* **77** 1413 (1996).
- [7] M. HORODECKI, P. HORODECKI AND R. HORODECKI, *Phys. Lett. A* **223** 1 (1996).
- [8] V. VEDRAL, M. B. PLENIO, M. A. RIPPIN AND P. L. KNIGHT, *Phys. Rev. Lett.* **78**, 2275 (1995).
- [9] V. VEDRAL AND M. B. PLENIO, *Phys. Rev. A* **57**, 1619 (1998).
- [10] W. K. WOOTTERS, *Phys. Rev. Lett.* **80** 2245 (1998).
- [11] M. LEWENSTEIN AND A. SANPERA, *Phys. Rev. Lett.* **80**, 2261 (1998).
- [12] B.-G ENGLERT AND N. METWALLY, *J. Mod. Opt.* **47**, 2221 (2000).
- [13] S. J. AKHTARSHENAS AND M. A. JAFARIZADEH, *eprint quant-ph/0207161* (2002).
- [14] F. VERSTRAETE, J. DEHAENE AND B. DEMOOR, *Phys. Rev. A* **64**, 010101 (2001).
- [15] F. VERSTRAETE, J. DEHAENE AND B. DEMOOR, *Phys. Rev. A* **65**, 032308 (2002).